

# New synonyms of two Arabian ants of the genus *Monomorium* Mayr, 1855 (Hymenoptera, Formicidae)

Mostafa R. Sharaf<sup>1</sup>, Cedric A. Collingwood<sup>2</sup>, Hathal M. Al Dhafer<sup>1</sup>,  
Mohammed S. Al mutairi<sup>3</sup>, Abdulrahman S. Aldawood<sup>1</sup>

**1** Plant Protection Department, College of Food and Agriculture Sciences, King Saud University, Riyadh 11451, P. O. Box 2460, Saudi Arabia **2** 18 Milton Street, Skipton, North Yorkshire, BD23 2ED, U. K.  
**3** Saudi Wildlife Authority, Riyadh 11575, P.O. Box 61681, Saudi Arabia.

Corresponding author: Mostafa R. Sharaf ([antsharaf@gmail.com](mailto:antsharaf@gmail.com))

---

Academic editor: M. Borowiec | Received 24 February 2015 | Accepted 14 May 2015 | Published 21 May 2015

---

<http://zoobank.org/3E64E35D-C57F-4D5C-A66A-AF8DDA0C710A>

---

**Citation:** Sharaf MR, Collingwood CA, Al Dhafer HM, Al mutairi MS, Aldawood AS (2015) New synonyms of two Arabian ants of the genus *Monomorium* Mayr, 1855 (Hymenoptera, Formicidae). ZooKeys 505: 51–58. doi: 10.3897/zookeys.505.9441

---

## Abstract

Synonymy of two Arabian *Monomorium* Mayr, 1855 species is proposed: *M. exiguum* Forel, 1894 = *M. desertorum* Collingwood & Agosti, 1996, **syn. n.**; *M. subopacum* Smith, 1858 = *M. mintiribe* Collingwood & Agosti, 1996, **syn. n.** A lectotype for *M. venustum* Smith, 1858 is designated. Information on nesting habits of *M. exiguum* and *M. venustum* in the Kingdom of Saudi Arabia are provided for the first time. Recently collected records for *M. exiguum*, *M. subopacum*, and *M. venustum* from the Kingdom of Saudi Arabia and United Arab Emirates are listed.

## Keywords

Arabian Peninsula, Middle East, Saudi Arabia, United Arab Emirates, synonymy, taxonomy, new designation

## Introduction

The first published work on the ant genus *Monomorium* Mayr, 1855 for the Kingdom of Saudi Arabia (KSA) was by Collingwood (1985), who listed and keyed 20 species from the country. The genus was subsequently treated comprehensively for the Arabian

Peninsula by Collingwood and Agosti (1996). The authors reported 53 species, 32 of which were described as new, (including 15 from the KSA, 10 from Oman, five from Yemen, and two from Kuwait). Collingwood et al. (2011) treated the myrmecofauna of the United Arab Emirates (UAE) and reported 29 *Monomorium* species. Three species were recorded from Socotra Island (Collingwood et al. 2004) and a new species, *M. nimihil* Collingwood, 2004 was described.

Recently the *Monomorium* fauna of KSA has received renewed attention, with the first record of *M. exiguum* Forel, 1894 (Aldawood and Sharaf 2009) and descriptions of three new species, *M. moathi* Sharaf & Collingwood, 2010 (Aldawood et al. 2010); *M. dryhimi* Aldawood & Sharaf, 2011 (Aldawood and Sharaf 2011) and *M. sarawatensis* Sharaf & Aldawood, 2013 (in El-Hawagry et al. 2013). During two visits to the World Museum, Liverpool, United Kingdom, two new synonyms were discovered for Arabian *Monomorium*.

## Materials and methods

### Abbreviations of museums

<b>BMNH</b>	The Natural History Museum, London, United Kingdom.
<b>MHNG</b>	Museum d'Histoire Naturelle, Geneva, Switzerland.
<b>NHMB</b>	Naturhistorisches Museum, Basel, Switzerland.
<b>WMLC</b>	World Museum Liverpool, Liverpool, United Kingdom.

The numbers between parentheses in material examined refer to individual workers.

## Results and discussion

### *Monomorium exiguum* Forel, 1894


Figs 1–4

For full synonymy see Heterick (2006), pp. 115–116.

*Monomorium exiguum* Forel, 1894: 85. (lectotype worker) Ethiopia. Afrotropic. “Ethiopia, Sudabessinien.” (MHNG), <http://www.antweb.org/specimen/CASENT0101870> [Image of type specimen examined].

*Monomorium desertorum* Collingwood & Agosti, 1996: 344 (w.) Saudi Arabia. Afrotropic.” (WMLC), <http://www.antweb.org/specimen/CASENT0906343>. Syntype worker [examined], Syn. n.

**Material examined.** Saudi Arabia, Baha, Dhi Ayn Archeological Village, 18.v.2010, 20.132°, 41.004°, 741m, (M. R. Sharaf, leg.) (21); Saudi Arabia, Riyadh, Oyaina, 28.iv.2010; 25.011°, 46.493°, 749m, (M. R. Sharaf, leg.) (3); Saudi Arabia, Riyadh, Qarina Valley, 5.xi.2009, 25.273°, 46.289°, 761m, (M. R. Sharaf, leg.) (3); Saudi Arabia, Baha,



**Figures 1–4.** *Monomorium exiguum* (worker), CASENT0101870. **1** Body in profile **2** Body in dorsal view **3** Head in full-face view **4** specimen label. Photo Zach Lieberman, <http://antweb.org/>

Dhi Ayn Archeological Village, 20.ix.2011, 20.132°, 41.004°, 744m, (M. R. Sharaf, leg.) (10); Saudi Arabia, Almajardah, wadi Khat, 10.xi.2012, 19.001°, 41.016°, 513m, (M. R. Sharaf, leg.) (6); Saudi Arabia, wadi Shahdan (Jizan), 13.xi.2012, 17.472°, 42.856°, 200m, (M. R. Sharaf, leg.) (8); Saudi Arabia, Wadi Aljora near Abadan, 12.xi.2012, 17.005°, 43.001°, 465m, (M. R. Sharaf, leg.) (6); Saudi Arabia, Baha, Wadi Elzaraeb, 9.v.2011, 20.073°, 41.387°, 2086m, (M. R. Sharaf, leg.) (1); Saudi Arabia, Abu Arish, 10.iv.2012, 17.013°, 42.802°, 90m, (M. R. Sharaf, leg.) (6); Saudi Arabia, Dhi Ayn Archeological Village, 11.v.2011, 19.929°, 41.442°, 741m, (M. R. Sharaf, leg.) (3); Saudi Arabia, Baha, Dhi Ayn Archeological Village, 7.iv.2013, 19.929°, 41.442°, 744m, (M. R. Sharaf, leg.) (4); Saudi Arabia, AlUrdiyah gov., Wadi Gonouna, 12.v.2011, 19.429°, 41.605°, 353m, (M. R. Sharaf, leg.) (20); Saudi Arabia, Al Bahah, Wadi Turabah, AlMandaq, 14.v.2011, 20.211°, 41.288°, 1793m, (M. R. Sharaf, leg.) (7); Saudi Arabia, Dhi Ayn Archeological Village, 15.v.2011, 19.929°, 41.442°, 741m, (M. R. Sharaf, leg.) (1); Saudi Arabia, Al Bahah, Wadi Turabah, AlMandaq, 10.v.2011, 20.211°, 41.288°, 1793m, (M. R. Sharaf, leg.) (1); Saudi Arabia, Riyadh, Hawtet Bani Tamim, 20.i.2014, 23.480°, 46.844°, 597m, (M. R. Sharaf, leg.) (3); Saudi Arabia, Al Qatif, El Naft, 23.iii.2012, 26.510°, 49.969°, 30m, (M. R. Sharaf, leg.) (2); UAE, Khor al-Khwair, 25.57.56.03, 8.iii.2007, (M. Hauser et al.) (1); UAE, Sharjah, 25.21.55.24, 28.ii–12.iv.2011, (M. Hauser et al.) (1); UAE, Wadi Bih dam, 25.48.56.04, 16–31.xii.2009, (M. Hauser et al.) (1).

**Remarks.** Only a single paratype specimen with the same data as the holotype exists at WMLC. The holotype and other paratypes are considered lost.

The description of *M. desertorum* in Collingwood and Agosti's (1996) indicated that the eyes are located anterior to the midlength of head, the scapes when retracted back do not reach the posterior margin of head, the antennae are 11 segmented, and the body is not sculptured except for the metanotal cross-ribs. Comparison was made of the single available paratype worker of *M. desertorum* with the lectotype worker of *M. exiguum* was carried out. We here treat *M. desertorum* as a junior subjective synonym of *M. exiguum*.

**Habitat.** The vast majority of *M. exiguum* nests that were collected in KSA were found to be associated with leaf litter and topsoil layers where workers foraged. Frequently nests were directly in the soil. The nesting habits of *M. exiguum* however, are diverse. In a site located in the southwestern mountains of the KSA, the species was found nesting in loose sandy soil with high moisture content and among roots of small *Portulaca oleracea* L. (Portulacaceae) plants beneath a date palm tree, *Phoenix dactylifera* L. (Arecaceae). Several worker series were nesting in a humid clay soil under banana trees. Other worker series were collected under a rock next to *Juniperus procera* Hochst. ex Endlicher (Cupressaceae) and *Acacia* spp. (Mimosaceae) trees. Another nest was found in thick layer of leaf litter under a large and old *Ficus benghalensis* L. (Moraceae) tree where the soil was rich in decayed organic matter. Some nests were found in leaf litter under *Calotropis procera* (Aiton) W.T.Aiton (Asclepiadaceae) and next to a mango tree (*Mangifera* sp., Anacardiaceae).

### ***Monomorium subopacum* (Smith, 1858)**

Figs 5–8

For full synonymy, see Heterick (2006), p. 103.


*Myrmica subopaca* Smith, 1858: 127 (w.q.) (paralectotype worker, designated by B. E. Heterick, September, 2004) Portugal (Madeira Is.). Afrotropic. "Portugal (Madeira Island), coll. T.V. Wollaston. (BMNH), <http://www.antweb.org/specimen/CASENT0010949> [Image of type specimen examined].

*Monomorium mintiribe* Collingwood & Agosti, 1996: 350, fig. 23 (w.q.m.) Oman. Palearctic. Bilad Ban. 17.i.1986, coll. W. Buttiker. (WMLC), Paratype worker [examined]. Syn. n.

**Material examined.** UAE, Ar-Rafah, 25.43.55.52, 1–8.iii.2011, (M. Hauser et al.) (1); UAE, Ar-Rafah, 25.43.55.52, 1.ii–31.iii.2010, (M. Hauser et al.) (1); UAE, Ar-Rafah, 25.18.56.07, 22.vi–2.vii.2010, (M. Hauser et al.) (1); UAE, Jebel Jibir, 25.39.56.07, 11–13.iv.2011, (M. Hauser et al.) (1).

**Remarks.** The holotype and 10 paratypes of *M. mintiribe* seem to be lost. Extensive searches at both WMLC and NHMB failed to locate type material except for a single paratype specimen labeled in red at the WMLC. In addition, despite the fact that the label information for the paratype specimen in WMLC does not exactly match the in-






**Figures 5–8.** *Monomorium subopacum* (paralectotype worker), CAsENT0010949. **5** Body in profile **6** Body in dorsal view **7** Head in full-face view **8** specimen label Photo April Nobile, <http://antweb.org/>

formation in Collingwood and Agosti (1996), we consider this specimen as part of the original type series. Collingwood and Agosti (1996) indicated the following paratypes: 1 male, 2 queens, 2 workers, “Oman, Bilad Bani, 20°03'N, 59°17'E, coll. W. Buttiker”, whereas the data on the paratype specimen in WMLC is “Bilad Ban, Oman, W. Buttiker, 17.i.1986.” The second author (C. A. Collingwood) confirms that the single remaining specimen is an originally designated paratype. The original description of *M. mintiribe* did not compare this taxon with related congeners. The single paratype is identical to the paralectotypes of *M. subopacum* and the original description agrees with this. Therefore, *M. mintiribe* is treated here as a junior subjective synonym of *M. subopacum*.

### *Monomorium venustum* (Smith, 1858)

Figs 9–12

*Myrmica venusta* Smith, 1858: 126 (w.) (lectotype worker) Syria. Palearctic. (BMNH “E” 1015257) [**new designation**].



**Figures 9–12.** *Monomorium venustum* (Lectotype worker), CASENT0902221. **9** Body in profile **10** Body in dorsal view **11** Head in full-face view **12** specimen label. Photo Will Ericson, <http://antweb.org/>

**Material examined.** Saudi Arabia, Al Atawla (Baha-Taif RD), Wadi Bawah, 8.xi.2012, 21.004°, 41.247°, 1310m, (M. R. Sharaf, leg.) (10); Saudi Arabia, Baha, Wadi Elzaraeb, 9.v.2011, 20.073°, 41.387°, 2086m, (M. R. Sharaf, leg.) (3); Saudi Arabia, Riyadh, Dirad, 30.xii.2009, 24.409°, 46.662°, 588m, (M. R. Sharaf, leg.) (6); Saudi Arabia, Al Bahah, Wadi Turabah, AlMandaq, 19.ix.2011, 20.242°, 41.262°, 1751m, (M. R. Sharaf, leg.) (6); Saudi Arabia, Riyadh, Alhota, 19.iv.2008, (M. R. Sharaf, leg.) (7); Saudi Arabia, Riyadh, Wadi Hanifa, 11.iv.2013, 24.671°, 46.595°, 641 (M. R. Sharaf, leg.) (14); Saudi Arabia, Al Bahah, Wadi Turabah, AlMandaq, 10.v.2011, 20.211°, 41.288°, 1751m, (M. R. Sharaf, leg.) (4).

**Remarks.** Originally, *M. venustum* was described based on syntypes of the worker caste from Syria. Here we designate a lectotype with the following data, “*M. venusta* Smith, type, BMNH (E), 1015257”. The Lectotype is deposited at BMNH.

**Habitat.** Workers of *M. venustum* build nests directly into the ground under stones and rocks, directly into the ground. This species apparently prefers to nest in soil with high moisture content as observed in many locations in KSA. In the southwestern mountains of the KSA, nests were constructed next to *Juniperus procera* Hochst. ex Endlicher (Cupressaceae) and *Acacia* spp. (Mimosaceae) trees. In addition, the species

is usually foraging in areas with dense green flowering grasses that covering the ground. A single nest was found existing next to *Mentha longifolia* (L.) Huds. (Lamiaceae). Myrmecophilous arthropods (e.g. small beetles, isopods and millipedes) were found inside some nests.

## Acknowledgment

The project was funded by the National Plan for Science, Technology and Innovation (MAARIFAH), King Abdulaziz City for Science and Technology, Kingdom of Saudi Arabia, award number 12-ENV2484-02. We are grateful to Prince Bandar Bin Saud Al Saud, the Head of the Saudi National Commission for Wildlife Conservation and Development for the generous support during the study. The authors are much grateful to the subject editor Marek Borowiec for his careful editing, Dr. Brian Fisher, Michele Esposito, Zach Lieberman, Estella Ortega, April Nobile, Ryan Perry and Will Ericson (California Academy of Sciences, San Francisco, USA) for photographing the type specimens of ants and making them accessible to researchers. We are indebted to Dr. Boris Kondratieff (Colorado State University) for the valuable suggestions and careful editing. Special thanks to Dr. Stephen Judd (Director, WML), Mr. Tony Hunter (Curator, Entomology, WML), and Isabelle Zürcher-Pfander (Collection manager, NHMB) for facilitating the study of the Arabian type materials. Mostafa Sharaf is particularly most grateful to the following: Suzanne Ryder (BMNH, London) for appreciated help to study type materials, Andrew Polaszek for his kindness and hospitality during the visit to the BMNH and finally to Wolfgang Schawaller and his wife Edith Schawaller for generous hospitality in Stuttgart.

## References

- Aldawood AS, Sharaf MR (2009) Two ant species (Hymenoptera: Formicidae) as new records to fauna of Saudi Arabia. *Bulletin of Entomological Society of Egypt* 86: 143–147.
- Aldawood AS, Sharaf MR, Collingwood CA (2010) *Monomorium moathi* sp. n., a new ant species from Yemen related to the *salomonis*-group. *Egyptian Academic Journal of Biological Sciences* 3(2): 37–42.
- Aldawood AS, Sharaf MR (2011) *Monomorium dryhimi* sp. n., a new ant species (Hymenoptera: Formicidae) of the *M. monomorium* group from Saudi Arabia, with a revised key to the Arabian species of the group. *ZooKeys* 106: 47–54. doi: 10.3897/zookeys.106.1390
- Collingwood CA (1985) Hymenoptera, Fam. Formicidae of Saudi Arabia. *Fauna of Saudi Arabia* 7: 230–301.
- Collingwood CA, Agosti D (1996) Formicidae of Saudi Arabia (part 2). *Fauna of Saudi Arabia* 15: 300–385.
- Collingwood CA, Agosti D, Sharaf MR, van Harten A (2011) Order Hymenoptera, Family Formicidae. *Arthropod Fauna of the UAE* 4: 405–474.

- Collingwood CA, Pohl H, Gusten R, Wranik W, van Harten A (2004) The ants of the Socotra Archipelago. *Fauna of Arabia* 20: 473–495.
- El-Hawagry MS, Khalil MW, Sharaf MR, Fadl HH, Aldawood AS (2013) A preliminary study on the insect fauna of Al-Baha Province, Saudi Arabia, with descriptions of two new species. *ZooKeys* 274: 1–88. doi: 10.3897/zookeys.274.4529
- Forel A (1894) Abessinische und andere afrikanische Ameisen, gesammelt von Herrn Ingenieur Alfred Ilg, von Herrn Dr. Liengme, von Herrn Pfarrer Missionar P. Berthoud, Herrn Dr. Arth. Müller etc. *Mitteilungen der Schweizerischen Entomologischen Gesellschaft* 9: 64–100.
- Heterick BE (2006) A revision of the Malagasy ants belonging to genus *Monomorium* Mayr, 1855. *Proceedings of the California Academy of Sciences* 57: 69–202.
- Smith F (1858) Catalogue of Hymenopterous insects in the collection of the British Museum (Part VI) – Formicidae. British Museum, London, 216 pp.

